

The Negative Relativistic Toda Hierarchy and Rational Poisson Brackets

Joana M. Nunes da Costa *

*Departamento de Matemática, Universidade de Coimbra, Apartado 3008,
3001-454 Coimbra, Portugal.
E-mail: jmcosta@mat.uc.pt*

and

Pantelis A. Damianou

*Department of Mathematics and Statistics, University of Cyprus, P. O. Box 20537,
1678 Nicosia, Cyprus.
E-mail: damianou@ucy.ac.cy*

We define hierarchies for negative values of the index of Poisson tensors, master symmetries and Hamiltonians for the finite, nonperiodic, relativistic Toda lattice. We show that the usual relations between master symmetries, Poisson tensors and invariants hold over all integer values of the index.

Key Words: Relativistic Toda lattice, bi-Hamiltonian system, Poisson brackets, master symmetries.

1. INTRODUCTION

The relativistic Toda lattice was introduced by S. N. Ruijsenaars [13] and was investigated in [1], [11], [12], [2] and [9]. In terms of canonical coordinates the relativistic Toda lattice is defined by the Hamiltonian

$$H(q^1, \dots, q^N, p_1, \dots, p_N) = \sum_{j=1}^N e^{p_j} f(q^{j-1} - q^j) f(q^j - q^{j+1}), \quad (1)$$

* This work was partially supported by CMUC-FCT

where $f(x) = \sqrt{1 + g^2 e^x}$ and, by convention, $q^0 = -\infty$, $q^{N+1} = \infty$. The number g is a coupling constant. To see the connection with the non-relativistic Toda lattice one writes the equation in Newtonian form

$$\ddot{q}^j = g^2 \dot{q}^j \left(\dot{q}^{j-1} \frac{e^{(q^{j-1} - q^j)}}{1 + g^2 e^{(q^{j-1} - q^j)}} - \dot{q}^{j+1} \frac{e^{(q^j - q^{j+1})}}{1 + g^2 e^{(q^j - q^{j+1})}} \right), \quad (2)$$

$j = 1, 2, \dots, N$. Setting $\dot{q}^j = \dot{Q}_j + c$ and letting $c \rightarrow \infty$ and $g \rightarrow 0$ in such a way that $gc = 1$, one obtains the equations of motion for the classical Toda lattice

$$\ddot{Q}_j = e^{Q_{j-1} - Q_j} - e^{Q_j - Q_{j+1}}. \quad (3)$$

In the classical case one uses a change of variables to prove integrability. We follow the same technique. Combining the changes of variables from [13], [1], we set

$$\begin{aligned} a_j &= g^2 e^{q^j - q^{j+1} + p_j} f(q^{j-1} - q^j) / f(q^j - q^{j+1}) \\ b_j &= \dot{q}^j - a_j \end{aligned} \quad (4)$$

In these variables the Hamiltonian is homogeneous quadratic and the equations of motion become

$$\begin{cases} \dot{a}_i = a_i(b_{i+1} - b_i + a_{i+1} - a_{i-1}) \\ \dot{b}_i = b_i(a_i - a_{i-1}), \end{cases} \quad (5)$$

where $i = 1, \dots, N$ and, by convention, $a_0 = a_N = 0$.

Following [2] we write these equations in Lax pair form (L, M) where

$$L = \begin{pmatrix} a_1 + b_1 & a_1 & 0 & \cdots & \cdots & 0 \\ a_2 + b_2 & a_2 + b_2 & a_2 & 0 & \cdots & 0 \\ a_3 + b_3 & a_3 + b_3 & a_3 + b_3 & a_3 & \cdots & 0 \\ \vdots & & \ddots & \ddots & & \vdots \\ a_{N-1} + b_{N-1} & \cdots & & \ddots & a_{N-1} + b_{N-1} & a_{N-1} \\ b_N & b_N & \cdots & \cdots & & b_N \end{pmatrix} \quad (6)$$

and

$$M = \begin{pmatrix} 0 & a_1 & 0 & \cdots & \cdots & 0 \\ 0 & -a_1 & a_2 & \cdots & & \vdots \\ 0 & 0 & -a_2 & a_3 & \cdots & \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & \ddots & \ddots & a_{N-1} \\ 0 & \cdots & \cdots & & 0 & -a_{N-1} \end{pmatrix}. \tag{7}$$

This shows that the functions $\bar{H}_{n-1} = \frac{1}{n} \text{Tr } L^n$ are constants of motion and therefore the system is integrable.

In the new coordinates a_j, b_j the symplectic bracket is transformed into a new quadratic Poisson bracket defined as follows:

$$\begin{aligned} \{a_i, a_{i+1}\} &= -a_i a_{i+1} \\ \{a_i, b_i\} &= a_i b_i \\ \{a_i, b_{i+1}\} &= -a_i b_{i+1}. \end{aligned} \tag{8}$$

All other brackets are zero. This bracket has $\det L = \prod_{i=1}^N b_i$ as Casimir, and the eigenvalues of L are in involution. We denote this bracket by π_1 . We next define a linear bracket π_0 as follows:

$$\begin{aligned} \{a_i, b_i\} &= a_i \\ \{a_i, b_{i+1}\} &= -a_i \\ \{b_i, b_{i+1}\} &= a_i. \end{aligned} \tag{9}$$

All other brackets are zero. In this bracket $\text{Tr } L$ is the Casimir and $\bar{H}_1 = \frac{1}{2} \text{Tr } L^2$ is the Hamiltonian. Therefore we have a bi-Hamiltonian system, a situation similar to the classical case. The bi-Hamiltonian formulation and the complete integrability of this system were established using different methods such as the Lax representation [2], master symmetries [11], [2] and recursion operators [1], [11]. Another approach to obtain the multi-Hamiltonian formulation for the relativistic Toda lattice, using Oevel’s theorem [10] and based on a method introduced by Das and Okubo [5] and Fernandes [6], was adopted in [9]. In that work a recursion operator is defined in the (q, p) space, the resulting sequence of Poisson tensors, master symmetries and invariants are then projected in the space of (a, b) variables to construct the usual multi-Hamiltonian hierarchies. The initial brackets of the hierarchy are π_0 and π_1 constructed above.

In this paper we extend to negative values of the index the usual hierarchies of Poisson tensors, master symmetries and Hamiltonians, for the finite nonperiodic relativistic Toda lattice. The extension is obtained in

natural (p, q) coordinates and also in (a, b) Flaschka coordinates. This procedure was already adopted in [3] for the finite non-periodic Toda lattice and we mimic the techniques of that paper. As was pointed out in [3] the negative hierarchy was proven useful, for example, in establishing the bi-Hamiltonian formulation of the Bogoyavlensky–Toda systems [4].

The paper is divided into 2 parts. In the first part (section 2) we recall the main results on the finite, nonperiodic relativistic Toda lattice, including the well-known positive hierarchies of Poisson tensors, master symmetries and Hamiltonians. In section 3 we introduce the negative relativistic Toda hierarchy, in (p, q) and (a, b) coordinates, and show that the relations between master symmetries, Poisson tensors and Hamiltonians, known for the positive hierarchies, hold for all integers values of the index, in both coordinates. We close with some additional results and give, for small dimensions, examples of the rational brackets and master symmetries.

2. THE RELATIVISTIC TODA LATTICE

We recall a theorem due to Oevel [10] that gives a method of generating master symmetries for non-degenerate bi-Hamiltonian systems.

Let $(M, \Lambda_0, \Lambda_1)$ be a non-degenerate bi-Hamiltonian manifold with a recursion operator $\mathcal{R} = \Lambda_1 \Lambda_0^{-1}$ and let $Y_1 = \Lambda_0 \nabla H_1 = \Lambda_1 \nabla H_0$ be a bi-Hamiltonian vector field on M , where Λ_i denotes also the matrix of the Poisson tensor Λ_i . We denote by Y_i the Hamiltonian vector field $\Lambda_i \nabla H_0$.

THEOREM 1. *Suppose that the vector field X_0 is a conformal symmetry for both Λ_0, Λ_1 and H_0 . i.e., for some scalars α, β , and γ we have*

$$\mathcal{L}_{X_0} \Lambda_0 = \alpha \Lambda_0, \quad \mathcal{L}_{X_0} \Lambda_1 = \beta \Lambda_1, \quad X_0(H_0) = \gamma H_0. \quad (10)$$

Then, the vector fields $X_i = \mathcal{R}^i X_0$ are master symmetries, the tensors $\Lambda_i = \mathcal{R}^i \Lambda_0$ are Poisson and we have, for $i, j = 0, 1, 2, \dots$

- (a) $[X_i, X_j] = (\beta - \alpha)(j - i)X_{i+j};$
- (b) $[X_i, Y_j] = (\beta + \gamma + (\beta - \alpha)(j - 1))Y_{i+j};$
- (c) $\mathcal{L}_{X_i} \Lambda_j = (\beta + (\beta - \alpha)(j - i - 1))\Lambda_{i+j};$
- (d) $X_i(H_j) = (\lambda + (\beta - \alpha)(j + i))H_{i+j}.$

Let us now briefly recall some results on the relativistic Toda lattice following [2], [9], [12]. We consider \mathbb{R}^{2N} with coordinates

$$(q^1, \dots, q^N, p_1, \dots, p_N).$$

Let Λ_0 be the Poisson tensor given by

$$\Lambda_0 = \sum_{i=1}^N e^{-p_i} \frac{\partial}{\partial q^i} \wedge \left(\frac{\partial}{\partial p_i} + \sum_{j=i+1}^N \frac{\partial}{\partial q^j} \right) + \sum_{i=1}^{N-1} e^{q^i - q^{i+1} - p_{i+1}} \left(\left(\frac{\partial}{\partial p_i} + \frac{\partial}{\partial q^{i+1}} \right) \right)$$

$$\wedge \left(\frac{\partial}{\partial p_{i+1}} + \sum_{j=i+2}^N \frac{\partial}{\partial q^j} \right) - \frac{\partial}{\partial p_{i+1}} \wedge \sum_{j=i+2}^N \frac{\partial}{\partial q^j} \right), \quad (11)$$

and Λ_1 be the canonical Poisson tensor,

$$\Lambda_1 = \sum_{i=1}^N \frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}. \quad (12)$$

In [9] it was proved that these Poisson tensors are compatible and also that $X = \Lambda_0 \nabla H_1 = \Lambda_1 \nabla H_0$ is a bi-Hamiltonian vector field, with

$$H_0 = \sum_{i=1}^N \left(e^{q^i - q^{i+1} + p_i} + e^{p_i} \right), \quad (13)$$

and

$$H_1 = \sum_{i=1}^N \left(\frac{1}{2} \left(e^{q^i - q^{i+1} + p_i} + e^{p_i} \right)^2 + e^{q^{i-1} - q^i + p_{i-1}} \left(e^{q^i - q^{i+1} + p_i} + e^{p_i} \right) \right). \quad (14)$$

By convention, $q^0 = -\infty$ and $q^{N+1} = +\infty$.

Consider the Flaschka-type transformation $F : \mathbb{R}^{2N} \rightarrow \mathbb{R}^{2N-1}$ [12], [9],

$$F : (q^1, \dots, q^N, p_1, \dots, p_N) \mapsto (a_1, \dots, a_{N-1}, b_1, \dots, b_N), \quad (15)$$

where $a_i = e^{q^i - q^{i+1} + p_i}$ and $b_i = e^{p_i}$. These are new coordinates in \mathbb{R}^{2N-1} .

The Poisson tensors Λ_0 and Λ_1 project by F onto bivectors on \mathbb{R}^{2N-1} ,

$$\pi_0 = \sum_{i=1}^{N-1} a_i \left(\frac{\partial}{\partial a_i} \wedge \left(\frac{\partial}{\partial b_i} - \frac{\partial}{\partial b_{i+1}} \right) + \frac{\partial}{\partial b_i} \wedge \frac{\partial}{\partial b_{i+1}} \right) \quad (16)$$

and

$$\pi_1 = \sum_{i=1}^{N-1} a_i \frac{\partial}{\partial a_i} \wedge \left(-a_{i+1} \frac{\partial}{\partial a_{i+1}} + b_i \frac{\partial}{\partial b_i} - b_{i+1} \frac{\partial}{\partial b_{i+1}} \right). \quad (17)$$

These are the Poisson tensors that we defined earlier and they provide a bi-Hamiltonian structure for the relativistic Toda lattice, with Hamiltonians

$$\bar{H}_0 = \sum_{i=1}^N (a_i + b_i), \quad \bar{H}_1 = \sum_{i=1}^N (a_{i-1} (a_i + b_i) + \frac{1}{2} (a_i + b_i)^2), \quad (18)$$

where, by convention, $a_0 = a_N = 0$. For $i \geq 0$, $\bar{H}_i = \frac{1}{i+1} \text{tr} L^{i+1}$ are constants of motion. The Poisson bracket associated with π_0 , which is linear, is the one given by (9), while the Poisson bracket corresponding to π_1 is quadratic and is given by (8).

Coming back to the natural (p, q) coordinates, since Λ_0 (and also Λ_1) is non-degenerate, we can define a recursion operator $\mathcal{R} = \Lambda_1(\Lambda_0)^{-1}$. Once we have a recursion operator \mathcal{R} , an infinite sequence of pairwise compatible Poisson tensors on \mathbb{R}^{2N} , $\Lambda_i = \mathcal{R}^i \Lambda_0$, and an infinite sequence of Hamiltonians H_i , given by $dH_i = {}^t\mathcal{R}(dH_{i-1})$ are obtained.

We take

$$X_0 = \sum_{i=1}^N \frac{\partial}{\partial p_i}, \tag{19}$$

and compute

$$\mathcal{L}(X_0)\Lambda_0 = -\Lambda_0, \quad \mathcal{L}(X_0)\Lambda_1 = 0 \quad \text{and} \quad X_0(H_0) = H_0.$$

So Oevel's theorem can be applied with $\alpha = -1$, $\beta = 0$ and $\gamma = 1$ [9]. We quote the results for RTL, in natural (p, q) coordinates, in the following theorem.

THEOREM 2.

- (i) $\Lambda_i = \mathcal{R}^i \Lambda_0$ are Poisson tensors, $i \geq 0$;
- (ii) the functions H_i , $i \geq 0$, are in involution with respect to all Λ_j ;
- (iii) $X_i(H_j) = (1 + i + j)H_{i+j}$, $i, j \geq 0$;
- (iv) $\mathcal{L}_{X_i} \Lambda_j = (j - i - 1)\Lambda_{i+j}$, $i, j \geq 0$;
- (v) $[X_i, X_j] = (j - i)X_{i+j}$, $i, j \geq 0$;
- (vi) $\Lambda_j \nabla H_i = \Lambda_{j-1} \nabla H_{i+1}$, $i \geq 0, j \geq 1$.

The infinite sequence $(\Lambda_i), i \in \mathbb{N}_0$, of higher order Poisson tensors on \mathbb{R}^{2N} reduce, by F , to an infinite sequence $(\pi_i), i \in \mathbb{N}_0$, of pairwise compatible Poisson tensors on \mathbb{R}^{2N-1} . Moreover, the master symmetries $X_i = \mathcal{R}^i X_0$, $i \in \mathbb{N}$, are projectable vector fields [9]. We denote by \bar{X}_i the projected vector fields. A new version of theorem 2 in (a, b) coordinates holds:

THEOREM 3.

- (i) π_i are Poisson tensors, $i \geq 0$;
- (ii) The functions \bar{H}_i , $i \geq 0$, are in involution with respect to all π_j ;
- (iii) $\bar{X}_i(\bar{H}_j) = (1 + i + j)\bar{H}_{i+j}$, $i, j \geq 0$;
- (iv) $\mathcal{L}_{\bar{X}_i} \pi_j = (j - i - 1)\pi_{i+j}$, $i, j \geq 0$;
- (v) $[\bar{X}_i, \bar{X}_j] = (j - i)\bar{X}_{i+j}$, $i, j \geq 0$;

$$(vi) \pi_j \nabla \bar{H}_i = \pi_{j-1} \nabla \bar{H}_{i+1}, \quad i \geq 0, j \geq 1.$$

We call the hierarchies of Poisson tensors, master symmetries and Hamiltonians given by theorems 2 and 3, in (p, q) and (a, b) coordinates respectively, the *positive relativistic Toda hierarchies*.

3. THE NEGATIVE RTL HIERARCHY

Our aim is to show that the relations of theorems 2 and 3 hold for any integer value of the index. For this purpose we define the *negative relativistic Toda hierarchy*. When the tensors are projected into the (a, b) space we obtain rational Poisson brackets. We imitate the techniques of [3].

Let \mathcal{N} be the inverse of the positive recursion operator \mathcal{R} ,

$$\mathcal{N} = \mathcal{R}^{-1} = \Lambda_0(\Lambda_1)^{-1},$$

which is given in matrix form by $\mathcal{N} = \begin{pmatrix} A & B \\ C & A^T \end{pmatrix}$, where A^T stands for the transpose of A and the matrices $A = (a_{ij})$, $B = (b_{ij})$ and $C = (c_{ij})$, with $1 \leq i, j \leq N$, are defined by

$$\begin{cases} a_{ij} = 0, & j < i \\ a_{ii} = e^{-p_i} + e^{q^{i-1} - q^i - p_i} \\ a_{i+1,i} = e^{q^{i-1} - q^i - p_i} \\ a_{i+2,i} = a_{i+3,i} = \dots = a_{N,i} = e^{q^{i-1} - q^i - p_i} - e^{q^i - q^{i+1} - p_{i+1}}, \end{cases}$$

$$\begin{cases} b_{ij} = -b_{ji} \\ b_{i+1,i} = b_{i+2,i} = \dots = b_{N,i} = e^{-p_i} + e^{q^{i-1} - q^i - p_i}, \end{cases}$$

$$c_{i,i+1} = c_{i+1,i} = e^{q^i - q^{i+1} - p_{i+1}}$$

and all the others entries c_{ij} are zero.

Let us denote by K_0 the conformal symmetry X_0 ,

$$K_0 = \sum_{i=1}^N \frac{\partial}{\partial p_i} = X_0,$$

and by K_i the vector fields $K_i = \mathcal{N}^i K_0$, $i = 0, 1, 2, \dots$. These vector fields are master symmetries.

We next apply Oevel's theorem to the RTL, using the negative recursion operator \mathcal{N} and the conformal symmetry K_0 . Since

$$\mathcal{L}_{K_0} \Lambda_1 = 0, \quad \mathcal{L}_{K_0} \Lambda_0 = -\Lambda_0, \quad K_0(H_1) = 2H_1,$$

the constants are, in this case,

$$\alpha_{\mathcal{N}} = 0, \quad \beta_{\mathcal{N}} = -1, \quad \gamma_{\mathcal{N}} = 2.$$

We remark that, obviously, $\alpha_{\mathcal{N}} = \beta$ and $\beta_{\mathcal{N}} = \alpha$.

Using the convention $X_{-i} = K_i$, for $i \geq 0$, we compute

$$[X_{-i}, X_{-j}] = [K_i, K_j] = (\beta_{\mathcal{N}} - \alpha_{\mathcal{N}})(j - i)K_{i+j} = (i - j)X_{-i-j},$$

and letting $m = -i$ and $n = -j$, we obtain

$$[X_m, X_n] = (n - m)X_{m+n}, \quad m, n \in \mathbb{Z}^-. \quad (20)$$

The same relation holds in (a, b) -coordinates:

$$[\bar{X}_m, \bar{X}_n] = (n - m)\bar{X}_{m+n}, \quad m, n \in \mathbb{Z}^-. \quad (21)$$

Consider the following equalities (see [10]) which involve the recursion operator \mathcal{R} and its inverse \mathcal{N} ,

$$\mathcal{L}_{X_0}\mathcal{R} = \mathcal{L}_{X_0}(\Lambda_1\Lambda_0^{-1}) = (\beta - \alpha)\mathcal{R}, \quad (22)$$

$$\mathcal{L}_{X_0}\mathcal{N} = \mathcal{L}_{X_0}\mathcal{R}^{-1} = (\alpha - \beta)\mathcal{R}^{-1} = (\alpha - \beta)\mathcal{N}. \quad (23)$$

Using these relations we obtain, for $j \in \mathbb{N}$,

$$\mathcal{L}_{X_0}\mathcal{R}^j = j(\beta - \alpha)\mathcal{R}^j \quad \text{and} \quad \mathcal{L}_{X_0}\mathcal{N}^j = j(\alpha - \beta)\mathcal{N}^j. \quad (24)$$

Also, taking into account that \mathcal{R} and \mathcal{N} are torsionless tensors, we have, for any vector field X ,

$$\mathcal{L}_{N^i X}\mathcal{R}^j = N^i(\mathcal{L}_X\mathcal{R}^j), \quad i, j \in \mathbb{N}. \quad (25)$$

Relations (22)-(25) allow us to compute the bracket of two master symmetries $K_i = X_{-i}$ and X_j , one in the negative hierarchy and the second in the positive one [3]. The result is

$$[X_{-i}, X_j] = [\mathcal{N}^i X_0, \mathcal{R}^j X_0] = (j + i)(\beta - \alpha)X_{j-i}. \quad (26)$$

In the case of the RTL, with $\alpha = 1$ and $\beta = 0$, (26) turns to

$$[X_{-i}, X_j] = (j + i)X_{j-i}. \quad (27)$$

This relation also holds in (a, b) -coordinates and we conclude that (20) and (21) hold for any integer value of the index.

We can establish the following theorem.

THEOREM 4. *The conclusions of Theorems 2 and 3 hold for any integer value of the index.*

Proof. The conditions (i) and (ii) of theorem 2 are a direct consequence of Oevel’s theorem and the properties of the recursion operators \mathcal{R} and \mathcal{N} [8]. The map F being a Poisson morphism, (i) of theorem 3 follows from (i) of theorem 2. The proof of (ii) of theorem 3 is the same as the corresponding in the case of Toda system [3].

Regarding condition (iii) of theorem 2, the only case we have to verify is $X_{-j}(H_i)$, with $i, j > 0$. Consider $i \geq j$ (the other case is similar),

$$\begin{aligned} X_{-j}(H_i) &= \langle {}^t\mathcal{R}^i(dH_0), \mathcal{N}^j X_0 \rangle = \langle dH_0, \mathcal{R}^{i-j} \rangle \\ &= X_{i-j}(H_0) = (1 + i - j)H_{i-j}. \end{aligned}$$

To prove condition (iii) of theorem 3, involving the master symmetries \bar{X}_i , the arguments used in [3] in the case of the (classical) Toda lattice, can be applied for the RTL to prove that if λ is an eigenvalue of the matrix L , the equation $\bar{X}_n(\lambda) = \lambda^{n+1}$ holds for $n \in \mathbb{Z}$. With $\bar{H}_j = \frac{1}{j+1} \text{tr} L^{j+1}$, we obtain, for $n < 0, j \geq 0$ and $j \neq -1 - n$,

$$\begin{aligned} \bar{X}_n(\bar{H}_j) &= \frac{1}{j+1} \sum_{k=1}^N \bar{X}_n(\lambda_k^{j+1}) \\ &= (1 + n + j)\bar{H}_{n+j}; \end{aligned}$$

we conclude that

$$\bar{X}_n(\bar{H}_m) = (1 + m + n)\bar{H}_{n+m} \tag{28}$$

holds for any $n, m \in \mathbb{Z}$.

Let us now show that conditions (iv) of theorems 2 and 3 hold for negative indices. We set $W_j = \Lambda_{1-j}$ and for $i, j \geq 0$, we compute, using Oevel’s theorem,

$$\mathcal{L}_{X_{-i}}\Lambda_{-j} = \mathcal{L}_{K_i}W_{1+j} = (\beta_{\mathcal{N}} + (\beta_{\mathcal{N}} - \alpha_{\mathcal{N}})(j - i))W_{1+j+i}$$

which gives

$$\mathcal{L}_{X_{-i}}\Lambda_{-j} = (-1 + i - j)\Lambda_{-i-j}.$$

Letting $m = -i$ and $n = -j$, we obtain

$$\mathcal{L}_{X_m}\Lambda_n = (n - m - 1)\Lambda_{m+n}, \quad m, n \in \mathbb{Z}^-. \tag{29}$$

Finally, using the same technique as before, we can compute $\mathcal{L}_{X_{-i}}\Lambda_j$, $i, j \geq 0$, and we conclude that (29) holds for all $m, n \in \mathbb{Z}$.

Switching to (a, b) -coordinates, we obtain

$$\mathcal{L}_{\bar{X}_m}\pi_n = (n - m - 1)\pi_{m+n}, \quad m, n \in \mathbb{Z}. \tag{30}$$

The condition (v) of both theorems was already proved. Finally, the proofs of (vi) of theorems 2 and 3 are similar to those in the case of the Toda system, see [3]. ■

Remark 5. As it was remarked in [3], $\bar{H}_n = \frac{1}{n+1}trL^{n+1}$ is undefined for $n = -1$. We have $\bar{X}_{-1}(\bar{H}_0) = N$ and, more generally, $\bar{X}_{-n}(\bar{H}_{n-1}) = N$. If we define

$$\bar{X}_m(\bar{H}_{-1}) = \lim_{n \rightarrow -1} (\bar{X}_m(\bar{H}_n)),$$

then

$$\begin{aligned} \bar{X}_m(\bar{H}_{-1}) &= \lim_{n \rightarrow -1} ((1 + m + n)\bar{H}_{n+m}) \\ &= m\bar{H}_{m-1}. \end{aligned}$$

Let us now show that the master symmetries X_n , $n \in \mathbb{Z}$, can be obtained once the hierarchy of Poisson tensors is known. In other words, we will show that all X_n are Hamiltonian vector fields associated with the same function f , w.r.t. the Poisson tensors of the hierarchy.

Consider the function $f = \sum_{i=1}^N q^i$. Then $X_n(f) = 0$, for all $n \in \mathbb{Z}$. In fact,

$$X_0(f) = \sum_{i=1}^N \frac{\partial f}{\partial p_i} = 0$$

and since $X_1 = \mathcal{R}X_0$ is given by [9]

$$\begin{aligned} X_1 &= \sum_{i=1}^N \left((1 - i) \left(e^{q^i - q^{i+1} + p_i} + e^{p_i} \right) + \sum_{j=i+1}^N \left(e^{q^j - q^{j+1} + p_j} + e^{p_j} \right) \right) \frac{\partial}{\partial q^i} \\ &\quad + \sum_{i=1}^N \left(e^{p_i} + i e^{q^i - q^{i+1} + p_i} + (2 - i) e^{q^{i-1} - q^i + p_{i-1}} \right) \frac{\partial}{\partial p_i}, \end{aligned}$$

where, by convention, $q^0 = -\infty$ and $q^{n+1} = +\infty$, we compute

$$X_1(f) = \sum_{i=1}^n \left((1 - i) \left(e^{q^i - q^{i+1} + p_i} + e^{p_i} \right) + \sum_{j=i+1}^n \left(e^{q^j - q^{j+1} + p_j} + e^{p_j} \right) \right) = 0.$$

Similarly, one proves that $X_2(f) = 0$. By induction we prove that $X_n(f) = 0$, for all $n \in \mathbb{N}$: if $X_{n-1}(f) = 0$, then

$$\begin{aligned} X_n(f) &= \frac{1}{n-2}[X_1, X_{n-1}](f) \\ &= \frac{1}{n-2}(X_1(X_{n-1}(f)) - X_{n-1}(X_1(f))) \\ &= 0. \end{aligned}$$

For the negative hierarchy of master symmetries, the proof is similar, with $X_{-n} = K_n$. The vector field $X_{-1} = K_1$ is given by

$$\begin{aligned} X_{-1} &= \sum_{i=1}^N \left(\sum_{j=1}^{i-1} (e^{-p_j} + e^{q^{j-1}-q^j-p_j}) + (i-N)(e^{-p_i} + e^{q^{i-1}-q^i-p_i}) \right) \frac{\partial}{\partial q^i} \\ &+ \sum_{i=1}^N \left(e^{-p_i} + (N+1-i)e^{q^{i-1}-q^i-p_i} + (1+i-N)e^{q^i-q^{i+1}-p_{i+1}} \right) \frac{\partial}{\partial p_i}, \end{aligned} \quad (31)$$

and we compute

$$\begin{aligned} X_{-1}(f) &= \sum_{i=1}^N \left(\sum_{j=1}^{i-1} (e^{-p_j} + e^{q^{j-1}-q^j-p_j}) + (i-N)(e^{-p_i} + e^{q^{i-1}-q^i-p_i}) \right) \\ &= 0. \end{aligned}$$

Now, we remark that $X_0 = \sum_{i=1}^N \frac{\partial}{\partial p_i}$ is the Hamiltonian vector field of f with respect to Λ_1 , i.e. $X_0 = -[\Lambda_1, f]$, where $[\cdot, \cdot]$ stands for the Schouten bracket [7]. Next, we show that every master symmetry X_n is an Hamiltonian vector field of f .

PROPOSITION 6. *For any $n \in \mathbb{Z}$, $X_n = -[\Lambda_{n+1}, f]$.*

Proof. We proceed by induction, starting with the positive hierarchy ($n \in \mathbb{N}_0$). For $n = 0$, we compute

$$-[\Lambda_1, f] = - \sum_{i,j=1}^N \left[\frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}, q^j \right] = X_0.$$

If $X_{n-1} = -[\Lambda_n, f]$, then

$$X_n = \frac{1}{n-2}[X_1, X_{n-1}]$$

$$= -\frac{1}{n-2}[X_1, [\Lambda_n, f]].$$

Using the generalized Jacobi identity for the Schouten bracket [7], we obtain

$$\begin{aligned} X_n &= -\frac{1}{n-2}[f, [X_1, \Lambda_n]] \\ &= -\frac{1}{n-2}[\mathcal{L}_{X_1}\Lambda_n, f] \\ &= -[\Lambda_{n+1}, f]. \end{aligned}$$

If $n < 0$, the proof is similar. \blacksquare

Remark 7. In a similar way, it is easy to prove that the Hamiltonians H_i are also determined from the knowledge of the Poisson brackets associated with (Λ_i) , $n \in \mathbb{Z}$ and the function f .

Next we exhibit, in (a, b) -coordinates, some rational Poisson brackets and master symmetries of the negative relativistic Toda hierarchy. We present the case $N = 3$, in order to simplify the notation. The bracket corresponding to the Poisson tensor π_{-1} on \mathbb{R}^{2N-1} , which is the projection of Λ_{-1} (by F), is the following:

$$\begin{aligned} \{a_1, a_2\}_{-1} &= \frac{1}{b_1 b_2 b_3} a_1 a_2 (b_1 - b_2 + b_3) \\ \{a_1, b_1\}_{-1} &= \frac{1}{b_1 b_2 b_3} a_1 (b_2 b_3 + a_1 b_3) \\ \{a_1, b_2\}_{-1} &= \frac{1}{b_1 b_2 b_3} (a_1 a_2 (b_2 - b_1) - a_1 b_3 (a_1 + b_1)) \\ \{a_1, b_3\}_{-1} &= \frac{1}{b_1 b_2 b_3} (-a_1 a_2 b_3) \\ \{a_2, b_1\}_{-1} &= \frac{1}{b_1 b_2 b_3} a_1 a_2 b_1 \\ \{a_2, b_2\}_{-1} &= \frac{1}{b_1 b_2 b_3} (a_2 b_1 (a_2 + b_3) + a_1 a_2 (b_3 - b_2)) \\ \{a_2, b_3\}_{-1} &= \frac{1}{b_1 b_2 b_3} a_2 b_1 (-a_2 - b_2) \\ \{b_1, b_2\}_{-1} &= \frac{1}{b_1 b_2 b_3} (a_1 b_1 (a_2 + b_3) + a_1 b_3 (a_1 + b_2)) \\ \{b_2, b_3\}_{-1} &= \frac{1}{b_1 b_2 b_3} (a_2 b_1 (a_2 + b_3) + a_1 a_2 b_3 + a_2 b_1 b_2). \end{aligned}$$

The denominator $b_1 b_2 b_3$ is equal to $\det L$, where L is the matrix (6) of the Lax pair (L, M) . So the Poisson tensor π_{-1} is defined on the open dense set $\det L \neq 0$.

The vector field X_{-1} in (p, q) -coordinates is given by (30). The expression of X_{-2} in the same coordinates is quite complicated. The projections of X_{-1} and X_{-2} in (a, b) coordinates, for $N = 3$, are the following:

$$X_{-1} = \left(-\frac{2a_1}{b_1} + \frac{a_1}{b_2}\right) \frac{\partial}{\partial a_1} - \frac{a_2}{b_2} \frac{\partial}{\partial a_2} + \left(1 - \frac{a_1}{b_2}\right) \frac{\partial}{\partial b_1} + \left(1 + \frac{2a_1}{b_1}\right) \frac{\partial}{\partial b_2} + \left(1 + \frac{a_2}{b_2}\right) \frac{\partial}{\partial b_3}$$

and

$$X_{-2} = \sum_{i=1}^2 u_i \frac{\partial}{\partial a_i} + \sum_{i=1}^3 v_i \frac{\partial}{\partial b_i}$$

with

$$\begin{aligned} u_1 &= -\frac{2a_1}{b_1^2} + \frac{a_1}{b_2^2} - \frac{2a_1}{b_1 b_2} - \frac{2a_1^2}{b_1^2 b_2} + \frac{a_1^2}{b_1 b_2^2} + \left(\frac{a_1}{b_3} + \frac{a_1 a_2}{b_2 b_3}\right) \left(\frac{1}{b_2} - \frac{1}{b_1}\right) \\ u_2 &= -\frac{a_2}{b_2^2} + \frac{a_2}{b_1 b_2} \left(1 - \frac{a_1}{b_2}\right) - \frac{a_2}{b_2 b_3} \left(2 + \frac{a_2}{b_2}\right) - \frac{a_2}{b_1 b_3} \left(1 + \frac{a_1}{b_2}\right) \\ v_1 &= \frac{1}{b_1} - \frac{a_1}{b_2^2} - \frac{a_1^2}{b_1 b_2^2} - \frac{a_1}{b_2 b_3} - \frac{a_1 a_2}{b_2^2 b_3} \\ v_2 &= \frac{2a_1 a_2}{b_1 b_2 b_3} + \frac{a_1 + a_2}{b_1 b_3} + \frac{a_1}{b_1 b_2} + \frac{a_2}{b_2 b_3} + \frac{a_1^2}{b_1^2 b_2} + \frac{2a_1}{b_1^2} + \frac{1}{b_2} \left(1 + \frac{a_1}{b_1}\right)^2 \\ v_3 &= -\frac{a_2}{b_1 b_2} + \frac{a_1 a_2}{b_1 b_2^2} + \frac{a_2}{b_2^2} + \frac{1}{b_3} \left(1 + \frac{a_2}{b_2}\right)^2. \end{aligned}$$

The function $\det L = b_1 b_2 b_3$ is a Casimir of π_1 . On the other hand, $trL = H_0$ is a Casimir of π_0 . More generally, we can prove:

PROPOSITION 8. *The function trL^{1-n} is a Casimir of π_n , for all $n \in \mathbb{Z} \setminus \{1\}$.*

Proof. For $n = 2$, the technique of [2] (p. 5525) can be applied word for word to show that trL^{-1} is a Casimir of π_2 , i.e. $\pi_2 \nabla(trL^{-1}) = 0$. But $trL^{-1} = -H_{-2}$. So we get $\pi_2 \nabla H_{-2} = 0$.

By theorem 3,

$$0 = \pi_2 \nabla H_{-2} = \pi_3 \nabla H_{-3} = \dots$$

and the result is proved for all $n > 1$. For $n = 0$, as we already remarked, $trL = H_0$ is a Casimir of π_0 . So, $\pi_0 \nabla H_0 = 0$ and using again the Lenard relations of theorem 3, we obtain

$$0 = \pi_0 \nabla H_0 = \pi_{-1} \nabla H_1 = \pi_{-2} \nabla H_2 \dots$$

and the result is proved for $n < 1$. ■

REFERENCES

1. M. BRUSCHI AND O. RAGNISCO, *Recursion operator and Bäcklund transformations for the Ruijsenaars Toda lattice*, Phys. Lett. A **129** (1988), 21–25.
2. P. A. DAMIANOU, *Multiple Hamiltonian structures for Toda-type systems*, J. Math. Phys. **35** (1994), 5511–5541.
3. P. A. DAMIANOU, *The Negative Toda Hierarchy and Rational Poisson Brackets*, J. Geom. Phys. **45** (2003), 184–202.
4. P. A. DAMIANOU, *On the bi-Hamiltonian structure of Bogoyavlensky-Toda lattices*, Nonlinearity **17** (2004), 397–413.
5. A. DAS AND S. OKUBO, *A systematic study of the Toda lattice*, Ann. Physics **190** (1989), 215–232.
6. R. L. FERNANDES, *On the mastersymmetries and bi-Hamiltonian structure of the Toda lattice*, J. Phys. A **26** (1993), 3797–3803.
7. J.-L. KOSZUL, *Crochet de Schouten-Nijenhuis et cohomologie*, in Élie Cartan et les mathématiques d’aujourd’hui, Astérisque, numéro hors série (1985).
8. F. MAGRI AND C. MOROSI, *A geometrical characterisation of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds*, Quaderno S 19, Univ. de Milano, (1984).
9. J. M. NUNES DA COSTA AND C.-M. MARLE, *Master symmetries and bi-Hamiltonian structures for the relativistic Toda lattice*, J. Phys. A **30** (1997), 7551–7556.
10. W. OEVEL, *A geometrical approach to integrable systems admitting time dependent invariants*, in Topics in Soliton Theory and Exactly Solvable Nonlinear Equations, ed. M. Ablowitz, B. Fuchssteiner and M. Kruskal, World Scientific, Singapore (1987).
11. W. OEVEL, B. FUCHSSTEINER, H. ZHANG AND O. RAGNISCO, *Mastersymmetries, angle variables, and recursion operator of the relativistic Toda lattice*, J. Math. Phys. **30** (1989), 2664–2670.
12. YU. B. SURIS, *On the bi-Hamiltonian structure of Toda and relativistic Toda lattices*, Phys. Lett. A **180** (1993), 419–429.
13. S. N. RUIJSENAARS, *Relativistic Toda Systems*, Commun. Math. Phys. **133** (1990), 217–247.